



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE CYCLOTOMIC FUNCTION.*

By DR. L. E. DICKSON, The University of Chicago.

1. If $p^n=1$ and $p^m \neq 1$ ($m < n$), p is called a primitive n th root of unity. Let $Q_n(x)$ be the equation whose roots are the various n th primitive roots of unity without repetition. Let $n=\nu p^r$, ν not being divisible by the prime p . We first prove that

$$(1) \quad Q_n(x) = Q_\nu(x^{p^r}) \div Q_\nu(x^{p^{r-1}}).$$

To show that the division is exact, let ξ_1, \dots, ξ_ν be the distinct ν th roots of unity. Then $\xi_1^p, \dots, \xi_\nu^p$ differ only as to order from ξ_1, \dots, ξ_ν . Hence

$$Q_\nu(x^{p^r}) = \prod_{i=1}^{\nu} (x^{p^r} - \xi_i^p), \quad Q_\nu(x^{p^{r-1}}) = \prod_{i=1}^{\nu} (x^{p^{r-1}} - \xi_i).$$

Since $y - \xi$ divides $y^p - \xi^p$, the division (1) is exact. If the value x makes the quotient vanish, then $(x^{p^r})^m = \xi_i^m = 1$ if and only if m is a multiple of ν , while $x^{\nu p^{r-1}} \neq 1$; hence x is a primitive n th root of unity.

We employ (1) as a recursion formula to determine $Q_n(x)$. As a permanent notation, set $n = p_1^{r_1} p_2^{r_2} \dots p_s^{r_s}$, where p_1, \dots, p_s are distinct primes. Then

$$(2) \quad Q_n(x) = Q_\mu(x^{p_1^{r_1} p_2^{r_2}}) Q_\mu(x^{p_1^{r_1-1} p_2^{r_2-1}}) \div Q_\mu(x^{p_1^{r_1} p_2^{r_2-1}}) Q_\mu(x^{p_1^{r_1-1} p_2^{r_2}},$$

where $\mu = n \div p_1^{r_1} p_2^{r_2}$. In general, for $N = p_1^{r_1} p_2^{r_2} \dots p_\sigma^{r_\sigma}$, we get

$$(3) \quad Q_n(x) = \frac{Q_{n/N}(x^N) \prod Q_{n/N}(x^{N/p^i p^j}) \dots}{\prod Q_{n/N}(x^{N/p^i}) \prod Q_{n/N}(x^{N/p^i p^j p^k}) \dots},$$

in which i, j, \dots range from 1, \dots, σ . Now $Q_1(x) = x - 1$. For $\sigma = s$, $N = n$, and (3) becomes

$$(4) \quad Q_n(x) = \frac{(x^n - 1) \prod (x^{n/p^i p^j} - 1) \dots}{\prod (x^{n/p^i} - 1) \prod (x^{n/p^i p^j p^k} - 1) \dots},$$

where in the denominator the products extend over the combinations 1, 3, 5, \dots at a time of p_1, \dots, p_s ; in the numerator, 2, 4, \dots at a time.

Conversely, (1) follows from (4). The terms of (4) in which p_1 does not enter explicitly combine into $Q_\nu(x^{p_1^{r_1}})$; those in which p_1 enters explicitly combine into $1 \div Q_\nu(x^{p_1^{r_1-1}})$.

2. The usual proof* of (4) is essentially only a verification. Since $Q_d(x) = 0$ gives all the primitive d th roots of unity without repetition, we have

$$(5) \quad x^n - 1 = \prod Q_d(x), \quad x^{n/p_1} - 1 = \prod Q_\delta(x), \dots$$

where d ranges over all the divisors of n ; δ over those of n/p_1 . When the products (5) are substituted in the second member of (4), every Q cancels except Q_n . In fact, if $d = p_1^{a_1} \dots p_t^{a_t} p_{t+1}^{r_{t+1}} \dots p_s^{r_s}$, where $t > 0$ and each $a_i < r_i$, Q_d divides exactly $A = 1 + {}_tC_2 + {}_tC_4 + \dots$ terms of the numerator of (4) and exactly $B = {}_tC_1 + {}_tC_3 + \dots$ terms of the denominator, ${}_tC_k$ being the number of combinations of t things k at a time. But $A - B = (1-1)^t = 0$.

3. From equation (4) follows as a corollary the important formula

$$(6) \quad \phi(n) = n \left(1 - \frac{1}{p_1}\right) \left(1 - \frac{1}{p_2}\right) \dots \left(1 - \frac{1}{p_s}\right),$$

where $\phi(n)$ denotes the number of positive integers not greater than n and relatively prime to n . Indeed, if ρ is a primitive n th root of unity, ρ^m is likewise if and only if m is relatively prime to n . But the degree of (4) evidently equals the right member of (6).

It follows from (4) that the polynomial $Q_n(x)$ has integral coefficients.

There are various proofs of the theorem that $Q_n(x)$ is algebraically irreducible, i. e., can not be expressed as a product of polynomials in x with rational coefficients.†

4. Theorem. For an integer x , the greatest common divisor g of $Q_n(x)$ and $x^{n/p_1} - 1$ is 1 or p_1 . If $g = p_1$, Q_n is not divisible by p_1^2 unless $n = p_1 = 2$, $x \equiv 3 \pmod{4}$, whence $Q_n = x + 1$.

Dividing the first equation (5) by the second, we get

$$(7) \quad (x^{n/p_1})^{p_1-1} + (x^{n/p_1})^{p_1-2} + \dots + x^{n/p_1} + 1 = Q_n(x) \cdot P(x),$$

$P(x)$ being a polynomial in x with integral coefficients. When the left member of (7) is divided by $x^{n/p_1} - 1$, the remainder is 1 or p_1 . Hence $g = 1$ or p_1 .

Let $g = p_1$, so that $x^{n/p_1} - 1 = kp_1$, k an integer. Substituting $kp_1 + 1$ for x^{n/p_1} in (7), we obtain $p_1 + \frac{1}{2}p_1(p_1 - 1)kp_1 + \text{terms in } p_1^2$. This is not divisible by p_1^2 if $p_1 > 2$, nor if $p_1 = 2$ and k is even. If $p_1 = 2$ and $k = 2l + 1$, then $x^{n/2} = 4l + 3$, whence $n/2$ must be odd and $x \equiv 3 \pmod{4}$. Suppose that $n > 2$ and $n/2 = p^r p_3^{r_3} \dots p_s^{r_s} = m$ odd. Performing in (4) the divisions of the type $(x^{2a} - 1) \div (x^a - 1)$, we get

$$(8) \quad \frac{x^m + 1}{x^{m/p} + 1} \cdot \prod_{i=3}^s \frac{x^{m/p^i} + 1}{x^{m/p^i} + 1} \cdot \prod_{i,j=3}^s \frac{x^{m/p^i p^j} + 1}{x^{m/p^i p^j} + 1} \cdot \prod_{i,j,k} \frac{x^{m/p^i p^j p^k} + 1}{x^{m/p^i p^j p^k} + 1} \dots$$

*On the general principle of the inversion involved, see Dedekind, *Crelle*, Vol. 54 (1857), pp. 1-26; Dirichlet-Dedekind, *Zahlentheorie*, p. 382; Bachmann, *Kreisheilung*, 1872, pp. 8-11, 16, and *Zahlentheorie*, I, pp. 40-42.

†See Bachmann, *Kreisheilung* (Leipzig. Teubner, 1872), pp. 31-43.

Since the exponents are all odd, each fraction or its inverse equals $1+f$, f containing an even number of powers of x . Hence Q is odd (cf. §5).

5. Theorem. For $n=p_1^{r_1} \dots p_s^{r_s}$ and x an integer, $Q_n(x)$ is divisible by p_1 if and only if x belongs to the exponent $\nu=n/p_1^{r_1}$ modulo p_1 ; in the contrary case, $Q_n(x) \equiv 1 \pmod{p_1}$.

By Fermat's theorem, $x^{p_1} \equiv x \pmod{p_1}$. Hence by (1), $Q_n(x) \equiv 1 \pmod{p_1}$ unless $Q_\nu(x) \equiv 0$. Now $Q_\nu(x)$ divides algebraically the function

$$(x^\nu - 1) \div (x^{\nu/p^i} - 1) = (x^{\nu/p^i})^{p^i-1} + \dots + x^{\nu/p^i} + 1 \quad (1 \leq i \leq s).$$

Hence if $x^{\nu/p^i} \equiv 1 \pmod{p_1}$, there is an integer k such that $kQ_\nu(x) \equiv p_i \pmod{p_1}$; whence Q_ν is not congruent to 0 $\pmod{p_1}$. There remains the case in which x^{ν/p^i} is not congruent to 1 $\pmod{p_1}$ for $i=2, \dots, s$. If $x^\nu \equiv 1 \pmod{p_1}$, x belongs to the exponent ν modulo p_1 and $Q_\nu \equiv 0$; if $x^\nu - 1$ is not congruent to 0, its divisor Q_ν is not congruent to 0 $\pmod{p_1}$.

Example. For $n=2.3.7$, formula (8) gives

$$Q_{42}(x) = \frac{(x^{21} + 1)(x + 1)}{(x^7 + 1)(x^3 + 1)} = x^{12} + x^{11} - x^9 - x^8 + x^6 - x^4 - x^3 + x + 1.$$

Thus $Q_{42} \equiv 1 \pmod{2 \text{ or } 3}$; $Q_{42}(x) \equiv 1 \pmod{7}$ if $x \equiv 0, -1$ or $x^3 \equiv +1$; but $Q_{42}(x) \equiv 0 \pmod{7}$ if $x^2 - x + 1 \equiv 0 \pmod{7}$, i. e., if x belongs to the exponent $\nu=6$.

Corollary. No one of the prime factors of n except the greatest can divide $Q_n(x)$.

6. Theorem. If x is a positive integer >1 , $Q_n(x)$ has a prime factor not dividing $x^n - 1$ ($n < n$), except in the cases $n=2$, $x=2^k - 1$ ($k \geq 2$); and $n=6$, $x=2$.

If $n=p^r$, $Q=y^{p^{r-1}} + \dots + y + 1 > p$, where $y=x^{p^{r-1}}$, so that the theorem follows from §4. We suppose henceforth that $n=p_1^{r_1} \dots p_s^{r_s}$, $s \geq 2$.

In view of §1, $Q_n = A/B$, where

$$A = \frac{x^n - 1}{x^{n/p_1} - 1} \cdot \prod \frac{x^{n/p_i p_j} - 1}{x^{n/p_i p_j} - 1} \dots, \quad B = \prod \frac{x^{n/p_i} - 1}{x^{n/p_i p_i} - 1} \cdot \prod \frac{x^{n/p_i p_i p_k} - 1}{x^{n/p_i p_i p_k} - 1} \dots,$$

in which i, j, k, \dots run from 2 to s . Now $x^{a(k-1)} < (x^{ka} - 1) \div (x^a - 1) < 2x^{a(k-1)}$. Hence $Q_n > \alpha/\beta$, where α is the result of retaining only the first term of each division in A , β the result of taking twice the first term of each division in B . The number of factors 2 introduced in B is $s-1 C_1 + s-1 C_3 + \dots = 2^{s-2}$. The exponent of x in α/β is evidently the degree $\phi(n)$ of Q_n . Let $x^{n/p_1 p_2 \dots p_s} = y$. Hence

$$Q_n(x) > y^{(p_1-1) \dots (p_s-1) \div 2^{s-2}},$$

y an integer >1 . In view of §4-5, it suffices to prove that $Q_n > p_1$, the greatest of the primes p_i , the case $n=6$, $x=2$ being an exception. For $s \geq 2$, we have $p_1 \geq 5$, $y^{p_1-1} > 2p_1$, the latter being true for $y=2$. Hence

$$Q_n > (2p_1)^{(p_2-1)\dots(p_s-1)} \div 2^{2^{s-2}} > p_1,$$

since at least $s-2$ of the primes p_2, \dots, p_s exceed 2, so that the exponent is $\geq 2^{s-2}$. For $s=2$, we have $p_1 \geq 3$, $Q_n > \frac{1}{2}y^{(p_1-1)(p_2-1)} \geq p_1$ unless $p_1=3$, $y^{p_1-1}=2$, whence $p_2=2$, $y=2$, $n=6$, $x=2$.

Corollary. *If x is a positive integer >1 , x^n-1 has a prime factor not dividing x^m-1 ($m < n$), except in the cases $n=2$, $x=2^k-1$; $n=6$, $x=2$.*

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

Problems 219, 220 were also solved by L. E. Newcomb. No. 222 was also solved by A. H. Holmes.

223. Proposed by THEODORE L. DE LAND, Office of the Secretary of the Treasury, Washington, D.C.

An officer in the Treasury Department assigned three clerks to count a lot of silver dollars and when finished noted that there was an apparent difference in their efficiency; and, to determine the fact, gave to each a similar lot of the same amount to count, the only record made at the time being that *A* to count his lot alone, took three weeks longer, *B* took two weeks longer, and *C* took one week longer than it took for all working together to count the first lot. The best counter, on the record made, was given an efficiency mark of 93 on the scale of 100. What efficiency mark should, on the record, be given to each of the other two counters?

Solution by the PROPOSER.

Let x = the time for *A*, *B*, and *C* working together to finish one lot.

Then $x+3$ = the time for *A* to finish one lot working alone;

$x+2$ = the time for *B* to finish one lot working alone; and

$x+1$ = the time for *C* to finish one lot working alone.

Then $\frac{1}{x}$ = what *A*, *B*, and *C* can do in one week working together;

$\frac{1}{x+3}$ = what *A* can do in one week working alone;

$\frac{1}{x+2}$ = what *B* can do in one week working alone; and

$\frac{1}{x+1}$ = what *C* can do in one week working alone.

Equating like terms we have,

$$\frac{1}{x} = \frac{1}{x+3} + \frac{1}{x+2} + \frac{1}{x+1} \dots\dots (1).$$